

Cannabidiol: Science, Marketing, and Legal Perspectives

Jenny L. Wiley, Camille K. Gourdet, and Brian F. Thomas



RTI Press publication OP-0065-2004

RTI International is an independent, nonprofit research organization dedicated to improving the human condition. The RTI Press mission is to disseminate information about RTI research, analytic tools, and technical expertise to a national and international audience. RTI Press publications are peer-reviewed by at least two independent substantive experts and one or more Press editors.

Suggested Citation

Wiley, J. L., Gourdet, C. K., and Thomas, B. F. (2020). *Cannabidiol: Science, Marketing, and Legal Perspectives*. RTI Press Publication No. OP-0065-2004. Research Triangle Park, NC: RTI Press. <https://doi.org/10.3768/rtipress.2020.op.0065.2004>

This publication is part of the RTI Press Research Report series. Occasional Papers are scholarly essays on policy, methods, or other topics relevant to RTI areas of research or technical focus..

RTI International
3040 East Cornwallis Road
PO Box 12194
Research Triangle Park, NC
27709-2194 USA

Tel: +1.919.541.6000
E-mail: rtipress@rti.org
Website: www.rti.org

©2020 RTI International. RTI International is a registered trademark and a trade name of Research Triangle Institute. The RTI logo is a registered trademark of Research Triangle Institute.



This work is distributed under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 license (CC BY-NC-ND), a copy of which is available at <https://creativecommons.org/licenses/by-nc-nd/4.0/legalcode>

<https://doi.org/10.3768/rtipress.2020.op.0065.2004>

www.rti.org/rtipress

Contents

About the Authors	i
Acknowledgments	ii
Abstract	ii
Introduction	1
Overview: Chemicals in Cannabis Plants	1
Cannabidiol's Effects in the Body	4
State of the Science	5
Marketing Claims and Risk of Product Contamination	7
Regulation/Legal Status of CBD	9
Conclusions and Recommendations	10
References	11

About the Authors

Jenny L. Wiley, PhD, is Distinguished Fellow in the Center for Drug Discovery at RTI International.

Camille K. Gourdet, JD, MA, is a research public health analyst in the Youth, Violence Prevention and Community Justice program at RTI International.

Brian F. Thomas, PhD, is the senior director of analytical chemistry, discovery sciences, and pharmaceuticals at Canopy Growth Corporation; at the time of writing, he was a principal scientist in Analytical Chemistry and Pharmaceuticals in Discovery Sciences at RTI International.

RTI Press Associate Editor

Ina Wallace

Acknowledgments

Preparation of this manuscript supported by RTI International internal funds and by grants from the National Institute on Drug Abuse (DA-045003) and the National Center for Complementary and Integrative Health (AT-010773).

Abstract

Recent loosening of legal restrictions on cannabis and its chemical constituents, including phytocannabinoids such as Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD), has led to rapid proliferation and wide availability of products containing CBD. Although using pure CBD does not result in THC-like intoxication, it is not risk-free. In this review, we examine CBD from scientific, marketing, and regulatory perspectives. Specifically, we evaluate the evidence used to support statements concerning CBD's real and putative medical effects and discuss misleading information that has been used in marketing approaches. Also, we explore the current legal landscape surrounding CBD. We conclude that further research is necessary to clarify legitimate therapeutic effects of CBD. Federal regulation is also necessary to assure quality, safety, and efficacy of CBD products. Until new regulations are enacted to ensure purity and label accuracy, consumers should balance any perceived benefits of CBD use against potential risks associated with using products of unknown quality.

Introduction

Products that contain CBD (an abbreviation for cannabidiol) have become popular over the last couple of years, with easy availability not only in pharmacies, but also in convenience stores, veterinary offices, and online. Yet, widespread misconceptions about CBD's effects that are furthered by misleading advertisements leave consumers wondering about which sources of information to trust. The purpose of this paper is to provide consumers with a "layperson friendly" introduction to CBD. We begin by defining what CBD is and distinguishing it from other chemicals contained in the cannabis plant, followed by a brief overview of CBD's actions in the body. We continue with discussions of CBD from scientific, marketing, and regulatory perspectives and conclude with recommendations for cautious consumers. This paper will provide a framework for consumers to understand the rapidly expanding science and evolving regulations in this area.

Overview: Chemicals in Cannabis Plants

Cannabis sativa and *Cannabis indica* (botanical names of strains of the cannabis plant) contain myriad chemicals, including more than 500 identified cannabinoids and terpenes (Andre, Hausman, & Guerriero, 2016; Thomas & ElSohly, 2015). Although phytocannabinoids are found only in the cannabis plant, terpenes are essential oils contained in many types of plants and plant-based foods (e.g., broccoli, lemons, peppermint) and contribute to their characteristic flavors and aromas (Figure 1). Cannabinoids and terpenes are produced in resin-filled trichomes, small hair-like growths located most abundantly in the flowers of unpollinated female cannabis plants. Cannabinoids are believed to have a protective function for the plant, helping to defend the plant against insects, bacteria/fungi, and environmental stresses (Pate, 1994; Premoli et al., 2019). For humans, the plant has served many purposes over the course of history, being used for its medicinal and nutritional properties,

Figure 1. Terpene-containing plants and foods



Note: Besides cannabis, many other plants and foods contain terpenes, including those pictured here. Top row (l-r): *Cannabis sativa*, citrus fruits, spicy grass basil. Bottom row (l-r): eucalyptus leaves, cloves, beer hops.

in spiritual ceremonies, and as a source of fiber for rope and clothing (Andre et al., 2016; Zuardi, 2006). Cannabis also has a long history of being used for its intoxicating properties (Pollan, 2001).


Although cannabis contains numerous chemical constituents, two are key to current discussions of medicinal properties of the plant: Δ^9 -tetrahydrocannabinol (THC) and cannabidiol (CBD) (Figure 2). THC is primarily responsible for cannabis's intoxicating effects (reviewed in Gaoni & Mechoulam, 1964), although it may also contribute to therapeutic effects of the cannabis plant (Wilkinson et al., 2003). Through selective breeding and indoor cultivation, THC's concentration in cannabis strains used specifically for intoxication has increased dramatically in recent years (ElSohly et al., 2016). A flowering top (or "bud") of a *Sinsemilla* (female, nonfertilized) cannabis strain high in THC may be expected to contain levels of up to 20 to 25 percent total THC (primarily in its acid form); relatively simple processes can be used to concentrate the THC into hashish or hash oil preparations exceeding 50 percent total THC content (Chandra et al., 2019; Jikomes & Zoorob, 2018; Smart, Caulkins, Kilmer, Davenport, & Midgette, 2017). More complex extraction processes or purification procedures can produce cannabis extracts with THC concentrations of 68 percent or higher that are increasingly encountered in the consumer market (Hädener, Vieten, Weinmann, & Mahler, 2019; Raber, Elzinga,

& Kaplan, 2015; Smart et al., 2017; Stogner & Miller, 2015). Conversely, high-CBD strains may contain 10.9 to 18.9 percent CBD (primarily in its acid form), with minimal THC content (Rahn, 2018). Simple processing procedures also may be used to increase CBD concentrations in products. Unlike THC, CBD does not produce intoxication (Schoedel et al., 2018); however, it does reach the brain when ingested, smoked, or vaped at high enough concentrations (Alozie, Martin, Harris, & Dewey, 1980). Therefore, CBD may induce changes in functioning of the central nervous system.

Cannabis (and its extracts) marketed as "medical" may contain THC, CBD, or a combination of both, as well as other minor cannabinoids and terpenes. Cannabis marketed as "recreational" typically contains high concentrations of THC, often with very little CBD. The degree to which the blend of chemicals contained in a cannabis strain contribute to its pharmacological or therapeutic effects directly or through modulation of the activity of THC or CBD (i.e., the "entourage effect"; Russo, 2011) is an area of active research and has not yet been resolved.

In a freshly harvested cannabis plant, THC and CBD are present in their acidic forms (THCA and CBDA, respectively), and it is primarily when they are heated that the decarboxylation necessary for them to be converted to their more commonly known "free" (and pharmacologically active) forms occurs. Heating may occur during smoking/vaping or it may occur

Figure 2. Chemical structures of cannabidiol (CBD) and Δ^9 -tetrahydrocannabinol (THC)



Source: Adapted from Thomas and ElSohly (2015), Figure 1.1. © 2015, Elsevier in cooperation with RTI Press.

during the process of extraction from the plant, as in the creation of CBD oil. Although heating cannabis to high temperatures during extraction converts the acidic cannabinoids into active compounds, it also may remove many of the volatile terpenes (Romano & Hazekamp, 2013). Methods of extraction are variable, but the most common procedures rely on using a solvent to separate and concentrate the desired chemicals from the plant material. Solvents typically used for extraction include compressed gases (e.g., CO₂ in supercritical fluid extraction processes or butane) and other liquids (e.g., ethanol, olive oil). Ideally, following extraction, the resulting oil is purified to remove the solvent and any contaminants (e.g., heavy metals, pesticides) that may have been extracted from the plant along with the cannabinoids.

We will not discuss the intricacies of cannabinoid extraction and purification techniques, as they are beyond the scope of this paper. The important points to know about the extraction process are that (1) extraction aims to purify and concentrate (and may decarboxylate) the cannabinoids present in the plant material, resulting in higher cannabinoid amounts per unit volume than in the plant; (2) other undesired substances may also be extracted and concentrated during the process; and (3) in the absence of adequate purification, contaminants (including the solvent) may remain in the final product.

CBD may also be extracted from “hemp” plants. Although industrial hemp is a variant of *Cannabis sativa*, selective breeding has resulted in divergence between plants used for fiber and those strains consumed by humans. For example, hemp plants that are bred for industrial uses (e.g., fiber for textiles or rope) have a different appearance (e.g., they are taller and more fibrous stalks with thinner leaves) and typically do not have a high cannabinoid content (Figure 3). In contrast, “hemp” that is selectively bred for consumer use may have resin-rich flowers that contain high concentrations of CBD (acid form), and the flowers of cannabis plants bred for recreational use usually contain high concentrations of THC (acid form) (Andre et al., 2016; VanDolah, Bauer, & Mauck, 2019). In the United States, the classification of “hemp” is restricted legally to plants that contain less than 0.3 percent THC, although the CBD concentration of hemp is not controlled. However, if not carefully purified following extraction, hemp-derived CBD will invariably contain a modicum of THC due to concentration during extraction (see explanation in previous paragraph), which may affect its legality under federal law (Hilderbrand, 2018).

Hemp seeds (or uncontaminated oil from the seeds) do not contain CBD or THC but are rich in omega-3 and omega-6 polyunsaturated fatty acids in a nutritionally favorable 3:1 ratio (Siano et al., 2018). The US Food and Drug Administration (FDA), which

Figure 3. Comparison of “hemp” and “marijuana” strains of cannabis plants



Note: The industrial “hemp” strain of cannabis (panel A) is taller and has more fibrous stalks with thinner leaves. It is typically cultivated outdoors. The “marijuana” strain of cannabis (panel B) is characterized by its resin-rich flowers and is often cultivated indoors under strictly controlled light and temperature conditions.

is charged with ensuring the safety of the nation's food supply (among its other responsibilities), classifies hemp seed and oil as "generally recognized as safe" (GRAS) food additives (<https://www.fda.gov/food/cfsan-constituent-updates/fda-responds-three-gras-notice-hemp-seed-derived-ingredients-use-human-food>). Despite the recent upsurge of interest in CBD-infused foods, CBD does not share a GRAS designation as of this writing. Hence, consumers should not assume that consumption of CBD in a food or beverage product is recognized by the FDA as safe.

In the subsequent sections of this manuscript, we briefly describe the effects of CBD in the body and then examine CBD from scientific and marketing perspectives. We discuss the legal/regulatory status of CBD in the United States and conclude with recommendations for policy.

Cannabidiol's Effects in the Body

CBD is a highly lipid-soluble chemical with relatively low bioavailability (as low as 6 percent), especially when administered orally (Grotenhermen, 2003; Lucas, Galettis, & Schneider, 2018). Low bioavailability means either that low amounts of CBD are absorbed into the bloodstream or that large amounts of CBD are metabolized by the liver before reaching the presumed site of action (e.g., the brain or other body organ). To compensate for this rapid metabolism, an orally administered dose must be correspondingly increased. Among the factors that may affect bioavailability is the presence of food in the gastrointestinal tract, with high-fat meals being associated with greater CBD absorption (Birnbaum et al., 2019; Taylor, Gidal, Blakey, Tayo, & Morrison, 2018). CBD absorption into the plasma is also more efficient via smoking/vaping (estimated 31 percent bioavailability; [Agurell et al., 1986; Grotenhermen, 2003; Lucas et al., 2018; Ohlsson et al., 1986]), as this route of administration avoids exposure to gastric juices and first-pass metabolism by the liver and provides greater opportunity for the parent compound to reach its site of action. However, vaping and smoking CBD are associated with their own sets of disadvantages and risks, which several

other studies have described (Callaghan, Allebeck, & Sidorchuk, 2013; Kenne, Fischbein, Tan, & Banks, 2017).

After oral administration, CBD is extensively metabolized, primarily by enzymes in the liver (Jiang, Yamaori, Takeda, Yamamoto, & Watanabe, 2011). Drug-drug interactions may occur with CBD because other drugs (e.g., antiepileptics) that are metabolized by the same enzymes may compete with CBD for access to the enzymes (Gaston, Bebin, Cutter, Liu, & Szaflarski, 2017; Lucas et al., 2018). The most abundant CBD metabolite is 7-carboxy-CBD; lesser metabolites also form (Taylor et al., 2018). The degree to which these metabolites may have action(s) on their own has not been fully investigated. The primary route of elimination is through the feces (Grotenhermen, 2003; Lucas et al., 2018; Taylor et al., 2018). The time course of a single dose of oral CBD shows considerable variability within and among individuals, with a half-life ranging between 1 and 3 hours and time to maximal plasma level of approximately 4 to 5 hours (Millar et al., 2019; Taylor et al., 2018).

The mechanism by which CBD may produce any therapeutic effects is unclear. Although the chemical structure of CBD is somewhat similar to that of THC, CBD binds only poorly to CB₁ and CB₂ receptors ($K_i = 151$ and 4582 nM, respectively, with lower numbers indicating higher affinity) (Husni et al., 2014). These receptors are part of the body's endocannabinoid system, a system that is intimately involved in physiological processes related to the maintenance of homeostasis, including appetite regulation, energy balance, pain sensation, response to stress, and sleep (Di Marzo & Petrosino, 2007; Riebe & Wotjak, 2011; Rodríguez de Fonseca et al., 2005; Ruiz de Azua & Lutz, 2019).

CB₁ and CB₂ receptors are the two identified receptors in this system and are activated by cannabinoids such as anandamide and 2-arachidonoylglycerol that are naturally produced in the body and brain. Although CB₁ receptors are widely distributed in the brain and localized in several peripheral organs (e.g., lungs, liver), CB₂ receptors are primarily found in the periphery,

including the immune system (Galiègue et al., 1995; Kaminski, 1996). THC's direct activation of CB₁ receptors in the brain results in the characteristic subjective effects (e.g., intoxication) associated with the use of cannabis (Huestis et al., 2001; Wiley, Lowe, Balster, & Martin, 1995). CBD does not have a similar direct effect at CB₁ or CB₂ receptors, but it may affect the endocannabinoid system indirectly by attenuating anandamide metabolism, resulting in persistence of the endocannabinoid's interaction with the receptor (De Petrocellis et al., 2011). Unlike THC, however, receptor activation by endocannabinoid release in the brain does not result in psychoactive effects. Hence, CBD is not associated with such effects (Babalonis et al., 2017; Schoedel et al., 2018).

Based upon *in vitro* research, primarily in cellular models, CBD has also been reported to modulate activity of cannabinoid receptors without binding directly to the same molecular recognition site as THC or the endocannabinoids (i.e., allosteric modulation; Tham et al., 2019) and to act through other non-cannabinoid receptor systems, including transient receptor potential vanilloid 1 type (TRPV1) channels (Costa, Giagnoni, Franke, Trovato, & Colleoni, 2004; Sagredo, Ramos, Decio, Mechoulam, & Fernández-Ruiz, 2007), peroxisome proliferator-activated receptors (PPAR) gamma (Esposito et al., 2011; Hegde, Singh, Nagarkatti, & Nagarkatti, 2015; Hind, England, & O'Sullivan, 2016), adenosine (Carrier, Auchampach, & Hillard, 2006; Mijangos-Moreno, Poot-Aké, Arankowsky-Sandoval, & Murillo-Rodríguez, 2014), and serotonin (Hind et al., 2016; Rock et al., 2012; Sonogo, Gomes, Del Bel, & Guimaraes, 2016). Through these mechanisms or other as-yet-unidentified mechanisms, CBD produces its pharmacological effects. Few studies, however, have provided conclusive links between CBD's demonstrated therapeutic effects and verified pharmacological mechanisms underlying the effect (Ibeas Bih et al., 2015). Consequently, statements implying that CBD produces effects through its action on homeostatic processes mediated by the endocannabinoid system are misleading. CBD's effects on this system are indirect at best and, as yet, have been demonstrated primarily in molecular models.

State of the Science

Although it has been increasing, rigorous research on the therapeutic effects of cannabinoids in humans remains sparse. In part, research in the United States has been hampered by continued classification of cannabis constituents (including CBD) as Schedule I drugs. According to the DEA classification scheme, a Schedule I drug does not have any defined medical uses. Regulations controlling the transportation and use of Schedule I drugs are strict, even when such drugs are being used for research. Further, state-level legalization of cannabis does not negate the federal prohibition, because researchers working in institutions that receive federal funds are bound by the terms of federal law. However, rigorous research with cannabis constituents, including CBD, is crucial for decreasing false claims, revealing efficacy where it exists, and determining potentially harmful effects. Consumers need knowledge of costs and benefits of CBD use to make informed decisions about whether or not to use CBD. To this end, we summarize results of the scientific research that has examined pharmacological effects of CBD, emphasizing studies performed with human subjects.

Despite the overwhelming amount of misinformation that plagues the CBD market, the unsupported claims about CBD's therapeutic potential are not, by themselves, evidence that CBD does not have medicinal effects. Indeed, in June 2018, the FDA approved Epidiolex, a purified plant-based CBD extract, for use as a treatment for Dravet syndrome and Lennox–Gastaut syndrome, severe forms of childhood epilepsy (US Food and Drug Administration, 2018). Pursuant to this approval, the efficacy of Epidiolex for treatment of severe pediatric epilepsy was confirmed in several randomized placebo-controlled clinical trials and open-label studies (O'Connell, Gloss, & Devinsky, 2017; Szaflarski et al., 2018). Although the toxicity profile for Epidiolex was generally favorable, all drugs have some side effects, and Epidiolex is no exception. Common adverse effects reported after use of Epidiolex include drowsiness, diarrhea, and elevation of transaminases (i.e., liver enzymes) (Huestis et al., 2019; Rubin, 2018; White, 2019).

CBD may also interact with other drugs, which may result in the need to adjust dosing regimens (Gaston et al., 2017; Huestis et al., 2019; Lucas et al., 2018; Morrison, Crockett, Blakey, & Sommerville, 2019). In addition to Epidiolex, Sativex (nabiximols), a cannabis plant-based oral spray that contains a 1:1 ratio of THC:CBD, is approved in Canada, Australia and parts of Europe for treating conditions such as spasticity associated with multiple sclerosis (Collin, Davies, Mutiboko, & Ratcliffe, 2007; Giacoppo, Bramanti, & Mazzon, 2017). Dronabinol (Marinol), an FDA-approved synthetic form of THC, has also been available for years as a prescription medicine for conditions such as chemotherapy-induced nausea and vomiting and AIDS wasting syndrome.

Research with CBD in animal models suggests that there are other potential therapeutic effects, including analgesia, anti-anxiety effects, antipsychotic effects, neuroprotective effects, anti-inflammatory effects, and cancer treatment (Crippa, Guimarães, Campos, & Zuardi, 2018; García-Arencibia et al., 2007; Gomes et al., 2014; Pellati et al., 2018; Premoli et al., 2019; Ward, Ramirez, Neelakantan, & Walker, 2011). However, preclinical studies of CBD effects have been rather sporadic over the years, and support for these specific therapeutic effects of CBD is not a coherent and convincing body of work. In addition, demonstrating an effect in an animal model of a disease or disorder is only one of the initial steps in drug development. Many drug candidates that show promise preclinically are not effective in humans and/or have unacceptable adverse effects (Henderson, Kimmelman, Fergusson, Grimshaw, & Hackam, 2013; Pound & Ritskes-Hoitinga, 2018).

A handful of recent well-controlled clinical studies have examined the effects of CBD in humans for conditions other than epilepsy. For example, Hurd et al. (2019) evaluated the effects of acute and repeated administration of CBD in abstinent patients diagnosed with heroin use disorder. Patients were assigned randomly to treatment groups and received placebo or one of two doses (400 or 800 mg) of CBD. Neither the patient nor the direct care medical staff knew which substance the patient received. Results showed that patients who received CBD reported fewer symptoms of anxiety and less craving induced

by cues previously related to drug use, which were the two primary dependent measures. Although results of one study are not conclusive proof that CBD is effective in the treatment of heroin use disorder, the reduction of symptoms associated with relapse that was noted in this well-controlled study is promising. Similarly, Masataka (2019) reported anxiety reduction in another double-blind placebo-controlled study conducted in adolescents diagnosed with social anxiety disorder.

CBD has also been evaluated as an adjunct to regularly prescribed antipsychotics in patients with schizophrenia (McGuire et al., 2018). This randomized, double-blind, placebo-controlled clinical trial was conducted across multiple medical facilities. Patients in the CBD treatment group showed significant improvement on a scale measuring positive symptoms of psychosis, suggesting that it may be effective as an adjunctive therapy when administered with an antipsychotic. Its effectiveness as a stand-alone antipsychotic was not evaluated.

With the exception of a few published results of clinical investigation, the extant scientific literature does little to illuminate the issue of CBD efficacy for the many conditions it is purported to treat. Several retrospective chart reviews, case reports, and survey data suggest that CBD may have beneficial therapeutic effects in the treatment of some conditions, including anxiety, pain, colitis, and sleep disruptions (Bitencourt & Takahashi, 2018; Corroon & Phillips, 2018; Couch, Maudslay, Doleman, Lund, & O'Sullivan, 2018; Crippa et al., 2018; Shannon, Lewis, Lee, & Hughes, 2019); other reviews suggest that current evidence of CBD efficacy in the treatment of these disorders is only modest at best, and tenuous in many cases (Black et al., 2019; Kuhathasan et al., 2019; White, 2019). For example, many of the studies described in these reviews are compromised by their small sample sizes, inconsistent results across studies, experimental induction of condition (e.g., anxiety) versus treatment of an existing disorder, and/or lack of important controls (e.g., no placebo comparison, reliance on self-report data, no masking of treatment condition to avoid unintentional bias) (Kuhathasan et al., 2019; Mandolini et al., 2018; Millar et al., 2019; White, 2019). Notably, including a control group

that does not receive an active drug is necessary to evaluate any drug's effectiveness (Dobrilla & Scarpignato, 1994; Munnangi & Angus, 2019). This control is especially critical for eliminating the potential influence of placebo effects in causing any beneficial therapeutic effects observed in CBD trials; placebo effects tend to be marked with cannabis-derived medications (Gertsch, 2018).

In summary, additional research, larger scale trials, and replication studies are required to consider CBD an effective treatment for many of the various disorders in which its use is proposed (Kuhathasan et al., 2019; Lötsch, Weyer-Menkhoff, & Tegeder, 2018; Shannon et al., 2019; White, 2019). To date, controlled studies of CBD for therapeutic indications in humans (other than epilepsy; e.g., Devinsky et al., 2018, 2019) are rare, although this is changing rapidly with recent loosening of legal restrictions and societal acceptance. For example, ClinicalTrials.gov lists several clinical trials in various phases that propose examining CBD as a treatment for several conditions, including tremor, anxiety, pain, and substance abuse disorders (<https://clinicaltrials.gov/ct2/home>). Results from these trials and from other well-controlled research will distinguish between conditions for which CBD shows promise as an effective remedy and those for which it is merely a profitable elixir without true therapeutic benefit.

Marketing Claims and Risk of Product Contamination

CBD was originally isolated from cannabis in the 1940s by Roger Adams (Adams, Hunt, & Clark, 1940). Intensive consumer interest in the properties of CBD is relatively recent, however, especially when compared with the long and storied history of cannabis and the persistent fascination with the psychoactive effects of THC. In the last few years, the CBD industry has become increasingly lucrative and was valued at \$170 million in 2016; it is forecast to be valued at several billion dollars by 2023 (Corroon & Phillips, 2018). CBD products are widely available for purchase from both online and brick-and-mortar stores, including pharmacies, gas stations, convenience stores, and pet supply stores. Available

formulations include oils, lotions, and tinctures as well as “vape juice” (i.e., liquids for fillable electronic cigarettes). In addition, periodic media reports suggest that the food and beverage industry has become interested in CBD-infused products, ranging from beverages (e.g., tea, coffee, wine) to snack foods and fast-food cheeseburgers.

Given the diverse scope of these products, CBD may be described by product marketers as a dietary supplement, a food additive, a cosmetic ingredient, or a drug, each of which would be regulated differently by the FDA. A drug undergoes a rigorous FDA approval process, with required demonstration of its relative safety and efficacy for the treatment of condition(s) for which it is approved. CBD went through this process in 2018 when it was approved as a drug for severe pediatric epilepsy and is currently undergoing clinical trials for other conditions. While FDA drug approval focuses on efficacy, regulations governing dietary supplements, cosmetics and food/beverages focus primarily on post-market safety. An added complication in the case of CBD is its history as a Schedule I drug under the DEA scheme; this classification generally removes a chemical from the FDA's purview.

The lack of regulatory structure around the sale and legal status of CBD has contributed to the current proliferation in CBD products. CBD product manufacturers have taken advantage of the federal legal gray area regarding the manufacturing, sale, and distribution of these products. However, because CBD is now the active ingredient of an approved drug (i.e., Epidiolex, as described in the previous section), the FDA has started to use its authority over drug regulation to stop product promoters from making unsubstantiated claims about potential therapeutic efficacy of CBD-containing products (reviewed in Mead, 2019).

Marketers, however, can phrase their advertisements to remain within the emerging regulatory framework but also take advantage of the public's willingness to embrace CBD as a putative treatment for a variety of conditions, including anxiety, insomnia, pain, neuroprotection, and cancer. For example, marketing on websites may rely on consumer reviews rather

than direct claims by the seller (the latter of which could bring FDA censure). Further, company websites may only post positive reviews and omit any negative comments. Alternatively, CBD marketers may refer to other websites that promote CBD use for various conditions but do not sell it. Another equivocal marketing strategy is to conflate the results of scientific research that examined the effects of THC or of cannabis (which contains THC as well as CBD and other cannabinoids) with effects of CBD alone.

In addition to false or misleading claims of therapeutic efficacy, other issues related to sparse federal regulation of these products have arisen. For example, inaccurate labeling is not uncommon, with several studies reporting that the CBD concentration listed on the label of a commercially available product significantly diverged from the concentration determined by independent laboratories experienced in analyzing cannabinoid content (Bonn-Miller et al., 2017; Pavlovic et al., 2018; Vandrey et al., 2015; White, 2019). Another study found that CBD concentration was not consistent across batches of the same product purchased at different times, which conceivably could complicate dosage decisions for treatment of a chronic condition (White, 2019). Although some companies advertise that their products are laboratory-tested for CBD content, these claims often lack verification because CBD products are not currently listed in the United States Pharmacopeia, an annual index of drugs and products that meet defined quality standards for identity, strength, purity, and similar characteristics. Even for products that do contain reliable quantities of CBD, however, concentrations may not be high enough to produce the desired pharmacological effects, especially with oral consumption (see discussion of bioavailability in previous section).

This point also highlights issues that plague all cannabis products used for medical purposes: the most common sources of information about appropriate dosing and adverse effects do not include trained health professionals, and dosage is not determined empirically through scientific study. Rather, consumers consult dispensary or retail staff, friends, online sites, or media reports to determine

dosing. In states where a medical referral is needed to purchase medical cannabis, doctors write a recommendation that allows the patient to purchase products labeled “medical” in cannabis dispensaries, but dosage is not specified as it is for prescription drugs. When treatment with CBD does not achieve the desired results, the primary losses may be time and money; however, if a consumer chooses to use CBD to treat a serious or potentially terminal condition (e.g., cancer), the consequences may be correspondingly more severe, including progression of the disease or death.

A final consideration related to the absence of effective CBD regulation is product contamination. Contamination may occur when there is inadequate purification following CBD extraction. Residual solvent or its byproducts, including carcinogenic polycyclic aromatic hydrocarbons, are contaminants identified in CBD products that likely result from insufficient removal of the solvent extractor (Romano & Hazekamp, 2013; White, 2019). Because CBD is extracted from a plant, it is also susceptible to sources of unintentional contamination that may occur with plant-based products, including the introduction of pesticides, heavy metals, and microorganisms (e.g., bacteria, fungus) (Dryburgh et al., 2018; Lenton, Frank, Barratt, Potter, & Decorte, 2018; Roberts, 2019). In the absence of purification, the extraction process may result not only in both enhanced CBD content and increased concentration of pesticide and other contaminants. When pesticide contamination does occur, high heat used during the extraction process or during vaping may alter the chemical composition of the pesticide, which could change its toxicity profile—a problem that does not typically occur with pesticides and food crops. The legal/regulatory quagmire surrounding CBD at the federal level complicates adequate redress for these problems, although states are beginning to intervene with developing guidelines for pesticides and purification (Seltenrich, 2019). Finally, although rare, there have been documented instances of CBD products being contaminated with synthetic cannabinoids (i.e., extremely potent THC-like compounds produced in illicit laboratories) (Horth et al., 2018).

Regulation/Legal Status of CBD

In the United States, CBD's current legal status is in flux because of complex and conflicting laws and a multilayered system of federal and state regulation. In general, however, two threshold considerations govern the answer to the overall question, "Is CBD legal?": (1) whether the CBD substance or product is derived from "hemp" or from THC-containing "marijuana;" and (2) whether one is answering this question within the context of federal or state law (Haffajee, MacCoun, & Mello, 2018). This circuitous route to determining the legality of CBD is based on how federal and state laws have evolved. Typically, lawmaking authority operates in a hierarchy, such that the higher authority controls when the laws of two jurisdictions conflict (USCS Const. Art. VI, Cl. 2). Therefore, if both a federal and state law govern a situation, the federal law usually prevails (USCS Const. Art. VI, Cl. 2). Similarly, a conflict between a state and local law typically results in the state law prevailing (USCS Const. Art. VI, Cl. 2). We consider each of these two major questions in terms of possession and use of CBD by an individual user.

Under federal law, CBD products that are derived from "marijuana" are illegal to buy, use, or possess, unless an individual has a Schedule I license (O'Connor & Lietzan, 2019). However, CBD products that are derived from "hemp," defined under federal law as cannabis plants that contain less than 0.3 percent THC by dry weight, will be regulated differently in the near future, pursuant to the new authority created under the 2018 Farm Bill. Two things must happen before the cultivation, sale, and distribution of hemp-derived products will formally begin: (1) the USDA must finalize its regulations and guidelines that will oversee the cultivation of hemp and the manufacturing of hemp-derived products; and (2) any state or Indian tribes that wish to assert "primary regulatory authority" over the cultivation, manufacturing, sale, and distribution of hemp within their jurisdiction must draft these plans and submit them to the Secretary of Agriculture for approval (84 Fed. Reg. 58522 at 58523 [10/31/2019]).

Once fully implemented, the 2018 Farm Bill will legalize the cultivation of hemp and hemp-derived

products and will remove its constituents (including CBD) from the list of substances scheduled under the federal Controlled Substances Act (Agriculture Improvement Act of 2018, 115 P.L. 334, 132 Stat. 4490 [2018]). Implementation of the law requires several steps. For example, the US Department of Agriculture (USDA) must develop specific regulations to govern how hemp, as a crop, must be grown to comply with federal law. Until full implementation of the 2018 Farm Bill, the 2014 Farm Bill is the controlling law, meaning that only hemp-derived CBD products that comply with the cultivation of "industrial hemp" under a particular state's pilot research program may be sold under federal law (USDA, 2019). Because the DEA continues to classify both THC and CBD as Schedule I substances (US DEA, n.d.), hemp-derived CBD products remain illegal under federal law at the time of this writing, although this law has not been stringently enforced, even in states without laws that allow CBD use for medical purposes.

Under state law, the legality of CBD products varies widely depending on the laws governing medical or recreational cannabis in a specific state. To date, THC-containing medical cannabis has been legalized in 33 states and the District of Columbia, and 11 of those states and the District of Columbia have also legalized recreational cannabis. For the most part, these states' medical and recreational cannabis laws have focused on limiting the amount of THC within various cannabis-infused products and requiring certain disclosures that warn and inform consumers about the risks of ingesting these THC-containing products. Much less attention has been focused on limiting or even disclosing the amount of CBD contained within the products. Some medical or recreational cannabis states that do mention and specifically allow CBD products classify these CBD products as separate products with specific limits for CBD and THC and require a particular ratio of CBD to THC. For example, Washington State allows and regulates "high CBD compliant products."

Although most of the 17 US states that have not legalized either medical or recreational cannabis do allow either low-THC or CBD-only products (defined according to these states' laws), an important difference vis-à-vis these states and states that have

legalized medical or recreational cannabis is that the former have not created a comprehensive regulatory framework that formally regulates and enforces the states' laws with respect to the cultivation, manufacture, sale or distribution of these products. For example, many of these states do not formally require these low-THC or CBD-only products to be tested, labeled in a specific way, or registered within a seed-to-sale tracking system that enables the state to monitor a product at each stage of its production.

Federal and state laws dealing with cannabis, in general, and with CBD, in particular, vary widely (see Table 1). With respect to hemp-derived CBD products, the 2018 Farm Bill, once fully implemented, may resolve many of the conflicts between federal and state statutes concerning the legality of CBD; however, laws that deal with CBD derived from THC-containing cannabis will continue to conflict. Further, enforcement of cannabis laws requires analysis of confiscated material which, in turn, relies upon identification of chemicals contained in the material. Determining the source of CBD (i.e., "hemp" vs. "marijuana") is not possible with traditional methods of analysis. Instead, material with THC content under 0.3 percent is presumed to be derived from hemp, regardless of its actual source. The development of a comprehensive national policy on cannabis is needed

to resolve the many confusing issues surrounding regulation of cannabis and its constituents.

Conclusions and Recommendations

CBD is not an inert substance. Like any ingested chemical, it has the potential to alter the functioning of physiological systems, and preliminary evidence suggests that it does. Whether it does so in a manner that results in clinically significant improvement of the various disorders for which it is promoted is still an open question. Certainly, the evidence for CBD efficacy in treating defined forms of severe pediatric epilepsy is strong and offers hope for many families. Although only further research will provide conclusive evidence of other therapeutic effects of CBD, increased regulatory control of the CBD market is needed now, a situation that the FDA recognized in its sponsoring of a recent public hearing on "Scientific Data and Info About Cannabis Products." Indeed, the lack of regulatory structure around the sale and legal status of CBD has contributed to the current morass of product proliferation. Until new regulations are enacted to ensure purity and label accuracy, consumers should balance any presumed, but unproven, benefits against potential risks associated with using CBD products of unknown quality.

Table 1. Summary of CBD current legal status under federal and state law

	Recreational		Medical	
	Where is it legal?		Where is it legal?	
What is it?	Under federal law: Hemp-derived (containing < 0.3% THC, per the 2018 Farm Bill)	Under state law: Cannabis plant-derived, containing THC (currently 11 states + D.C., all 11 of which have <i>also</i> legalized medical cannabis)	Under federal law: Hemp-derived (containing < 0.3% THC, per the 2018 Farm Bill)	Under state law: Cannabis plant-derived, containing THC (currently 33 States +D.C. have legalized medical cannabis)
Hemp-derived CBD	Yes	Yes	Yes	Yes
THC-containing cannabis	No	Yes	No	Yes

References

- Adams, R., Hunt, M., & Clark, J. (1940). Structure of cannabidiol, a product isolated for the marihuana extract of Minnesota wild hemp. *Journal of the American Chemical Society*, *62*(1), 196–200. <https://doi.org/10.1021/ja01858a058>
- Agurell, S., Halldin, M., Lindgren, J. E., Ohlsson, A., Widman, M., Gillespie, H., & Hollister, L. (1986). Pharmacokinetics and metabolism of delta 1-tetrahydrocannabinol and other cannabinoids with emphasis on man. *Pharmacological Reviews*, *38*(1), 21–43.
- Alozie, S. O., Martin, B. R., Harris, L. S., & Dewey, W. L. (1980). 3H-delta 9-Tetrahydrocannabinol, 3H-cannabinol and 3H-cannabidiol: Penetration and regional distribution in rat brain. *Pharmacology, Biochemistry, and Behavior*, *12*(2), 217–221. [https://doi.org/10.1016/0091-3057\(80\)90359-7](https://doi.org/10.1016/0091-3057(80)90359-7)
- Andre, C. M., Hausman, J. F., & Guerriero, G. (2016). Cannabis sativa: The plant of the thousand and one molecules. *Frontiers in Plant Science*, *7*, 19. <https://doi.org/10.3389/fpls.2016.00019>
- Babalonis, S., Haney, M., Malcolm, R. J., Lofwall, M. R., Votaw, V. R., Sparenborg, S., & Walsh, S. L. (2017). Oral cannabidiol does not produce a signal for abuse liability in frequent marijuana smokers. *Drug and Alcohol Dependence*, *172*, 9–13. <https://doi.org/10.1016/j.drugalcdep.2016.11.030>
- Birnbaum, A. K., Karanam, A., Marino, S. E., Barkley, C. M., Remmel, R. P., Roslawski, M., ... Leppik, I. E. (2019). Food effect on pharmacokinetics of cannabidiol oral capsules in adult patients with refractory epilepsy. *Epilepsia*, *60*(8), 1586–1592. <https://doi.org/10.1111/epi.16093>
- Bitencourt, R. M., & Takahashi, R. N. (2018). Cannabidiol as a therapeutic alternative for post-traumatic stress disorder: From bench research to confirmation in human trials. *Frontiers in Neuroscience*, *12*, 502. <https://doi.org/10.3389/fnins.2018.00502>
- Black, N., Stockings, E., Campbell, G., Tran, L. T., Zagic, D., Hall, W. D., ... Degenhardt, L. (2019). Cannabinoids for the treatment of mental disorders and symptoms of mental disorders: A systematic review and meta-analysis. *The Lancet. Psychiatry*, *6*(12), 995–1010. [https://doi.org/10.1016/S2215-0366\(19\)30401-8](https://doi.org/10.1016/S2215-0366(19)30401-8)
- Bonn-Miller, M. O., Loflin, M. J. E., Thomas, B. F., Marcu, J. P., Hyke, T., & Vandrey, R. (2017). Labeling accuracy of cannabidiol extracts sold online. *Journal of the American Medical Association*, *318*(17), 1708–1709. <https://doi.org/10.1001/jama.2017.11909>
- Callaghan, R. C., Allebeck, P., & Sidorchuk, A. (2013). Marijuana use and risk of lung cancer: A 40-year cohort study. *Cancer Causes & Control*, *24*(10), 1811–1820. <https://doi.org/10.1007/s10552-013-0259-0>
- Carrier, E. J., Auchampach, J. A., & Hillard, C. J. (2006). Inhibition of an equilibrative nucleoside transporter by cannabidiol: A mechanism of cannabinoid immunosuppression. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(20), 7895–7900. <https://doi.org/10.1073/pnas.0511232103>
- Chandra, S., Radwan, M. M., Majumdar, C. G., Church, J. C., Freeman, T. P., & ElSohly, M. A. (2019). New trends in cannabis potency in USA and Europe during the last decade (2008–2017). *European Archives of Psychiatry and Clinical Neuroscience*, *269*(1), 5–15. <https://doi.org/10.1007/s00406-019-00983-5>
- Collin, C., Davies, P., Mutiboko, I. K., & Ratcliffe, S., & the Sativex Spasticity in MS Study Group. (2007). Randomized controlled trial of cannabis-based medicine in spasticity caused by multiple sclerosis. *European Journal of Neurology*, *14*(3), 290–296. <https://doi.org/10.1111/j.1468-1331.2006.01639.x>
- Corroon, J., & Phillips, J. A. (2018). A cross-sectional study of cannabidiol users. *Cannabis and Cannabinoid Research*, *3*(1), 152–161. <https://doi.org/10.1089/can.2018.0006>
- Costa, B., Giagnoni, G., Franke, C., Trovato, A. E., & Colleoni, M. (2004). Vanilloid TRPV1 receptor mediates the antihyperalgesic effect of the nonpsychoactive cannabinoid, cannabidiol, in a rat model of acute inflammation. *British Journal of Pharmacology*, *143*(2), 247–250. <https://doi.org/10.1038/sj.bjp.0705920>
- Couch, D. G., Maudslay, H., Doleman, B., Lund, J. N., & O'Sullivan, S. E. (2018). The use of cannabinoids in colitis: A systematic review and meta-analysis. *Inflammatory Bowel Diseases*, *24*(4), 680–697. <https://doi.org/10.1093/ibd/izy014>

- Crippa, J. A., Guimarães, F. S., Campos, A. C., & Zuardi, A. W. (2018). Translational investigation of the therapeutic potential of cannabidiol (CBD): Toward a new age. *Frontiers in Immunology*, *9*, 2009. <https://doi.org/10.3389/fimmu.2018.02009>
- De Petrocellis, L., Ligresti, A., Moriello, A. S., Allarà, M., Bisogno, T., Petrosino, S., ... Di Marzo, V. (2011). Effects of cannabinoids and cannabinoid-enriched Cannabis extracts on TRP channels and endocannabinoid metabolic enzymes. *British Journal of Pharmacology*, *163*(7), 1479–1494. <https://doi.org/10.1111/j.1476-5381.2010.01166.x>
- Devinsky, O., Nabhout, R., Miller, I., Laux, L., Zolnowska, M., Wright, S., & Roberts, C. (2019). Long-term cannabidiol treatment in patients with Dravet syndrome: An open-label extension trial. *Epilepsia*, *60*(2), 294–302. <https://doi.org/10.1111/epi.14628>
- Devinsky, O., Patel, A. D., Thiele, E. A., Wong, M. H., Appleton, R., Harden, C. L., ... Sommerville, K., & the GWPCARE1 Part A Study Group. (2018). Randomized, dose-ranging safety trial of cannabidiol in Dravet syndrome. *Neurology*, *90*(14), e1204–e1211. <https://doi.org/10.1212/WNL.0000000000005254>
- Di Marzo, V., & Petrosino, S. (2007). Endocannabinoids and the regulation of their levels in health and disease. *Current Opinion in Lipidology*, *18*(2), 129–140. <https://doi.org/10.1097/MOL.0b013e32803dbdec>
- Dobrilla, G., & Scarpignato, C. (1994). Placebo and placebo effect: Their impact on the evaluation of drug response in patients. *Digestive Diseases (Basel, Switzerland)*, *12*(6), 368–377. <https://doi.org/10.1159/000171471>
- Dryburgh, L. M., Bolan, N. S., Grof, C. P. L., Galettis, P., Schneider, J., Lucas, C. J., & Martin, J. H. (2018). Cannabis contaminants: Sources, distribution, human toxicity and pharmacologic effects. *British Journal of Clinical Pharmacology*, *84*(11), 2468–2476. <https://doi.org/10.1111/bcp.13695>
- ElSohly, M. A., Mehmedic, Z., Foster, S., Gon, C., Chandra, S., & Church, J. C. (2016). Changes in cannabis potency over the last 2 decades (1995–2014): Analysis of current data in the United States. *Biological Psychiatry*, *79*(7), 613–619. <https://doi.org/10.1016/j.biopsych.2016.01.004>
- Esposito, G., Scuderi, C., Valenza, M., Togna, G. I., Latina, V., De Filippis, D., ... Steardo, L. (2011). Cannabidiol reduces A β -induced neuroinflammation and promotes hippocampal neurogenesis through PPAR γ involvement. *PLoS One*, *6*(12), e28668. <https://doi.org/10.1371/journal.pone.0028668>
- Galiègue, S., Mary, S., Marchand, J., Dussossoy, D., Carrière, D., Carayon, P., ... Casellas, P. (1995). Expression of central and peripheral cannabinoid receptors in human immune tissues and leukocyte subpopulations. *European Journal of Biochemistry*, *232*(1), 54–61. <https://doi.org/10.1111/j.1432-1033.1995.tb20780.x>
- Gaoni, Y., & Mechoulam, R. (1964). Isolation, structure, and partial synthesis of an active constituent of hashish. *Journal of the American Chemical Society*, *86*(8), 1646–1647. <https://doi.org/10.1021/ja01062a046>
- García-Arencibia, M., González, S., de Lago, E., Ramos, J. A., Mechoulam, R., & Fernández-Ruiz, J. (2007). Evaluation of the neuroprotective effect of cannabinoids in a rat model of Parkinson's disease: Importance of antioxidant and cannabinoid receptor-independent properties. *Brain Research*, *1134*(1), 162–170. <https://doi.org/10.1016/j.brainres.2006.11.063>
- Gaston, T. E., Bebin, E. M., Cutter, G. R., Liu, Y., & Szaflarski, J. P., & the UAB CBD Program. (2017). Interactions between cannabidiol and commonly used antiepileptic drugs. *Epilepsia*, *58*(9), 1586–1592. <https://doi.org/10.1111/epi.13852>
- Gertsch, J. (2018). The intricate influence of the placebo effect on medical cannabis and cannabinoids. *Medical Cannabis and Cannabinoids*, *1*(1), 60–64. <https://doi.org/10.1159/000489291>
- Giacoppo, S., Bramanti, P., & Mazzon, E. (2017). Sativex in the management of multiple sclerosis-related spasticity: An overview of the last decade of clinical evaluation. *Multiple Sclerosis and Related Disorders*, *17*, 22–31. <https://doi.org/10.1016/j.msard.2017.06.015>
- Gomes, F. V., Issy, A. C., Ferreira, F. R., Viveros, M. P., Del Bel, E. A., & Guimarães, F. S. (2014). Cannabidiol attenuates sensorimotor gating disruption and molecular changes induced by chronic antagonism of NMDA receptors in mice. *The International Journal of Neuropsychopharmacology*, *18*(5), pyu04. <https://doi.org/10.1093/ijnp/pyu041>
- Grotenhermen, F. (2003). Pharmacokinetics and pharmacodynamics of cannabinoids. *Clinical Pharmacokinetics*, *42*(4), 327–360. <https://doi.org/10.2165/00003088-200342040-00003>

- Hädener, M., Vieten, S., Weinmann, W., & Mahler, H. (2019). A preliminary investigation of lung availability of cannabinoids by smoking marijuana or dabbing BHO and decarboxylation rate of THC- and CBD-acids. *Forensic Science International*, 295, 207–212. <https://doi.org/10.1016/j.forsciint.2018.12.021>
- Haffajee, R. L., MacCoun, R. J., & Mello, M. M. (2018). Behind schedule—Reconciling federal and state marijuana policy. *The New England Journal of Medicine*, 379(6), 501–504. <https://doi.org/10.1056/NEJMp1804408>
- Hegde, V. L., Singh, U. P., Nagarkatti, P. S., & Nagarkatti, M. (2015). Critical role of mast cells and peroxisome proliferator-activated receptor gamma in the induction of myeloid-derived suppressor cells by marijuana cannabidiol in vivo. *Journal of Immunology*, 194(11), 5211–5222. <https://doi.org/10.4049/jimmunol.1401844>
- Henderson, V. C., Kimmelman, J., Fergusson, D., Grimshaw, J. M., & Hackam, D. G. (2013). Threats to validity in the design and conduct of preclinical efficacy studies: A systematic review of guidelines for in vivo animal experiments. *PLoS Medicine*, 10(7), e1001489. <https://doi.org/10.1371/journal.pmed.1001489>
- Hilderbrand, R. L. (2018). Hemp & cannabidiol: What is a medicine? *Missouri Medicine*, 115(4), 306–309.
- Hind, W. H., England, T. J., & O’Sullivan, S. E. (2016). Cannabidiol protects an in vitro model of the blood-brain barrier from oxygen-glucose deprivation via PPAR γ and 5-HT $1A$ receptors. *British Journal of Pharmacology*, 173(5), 815–825. <https://doi.org/10.1111/bph.13368>
- Horth, R. Z., Crouch, B., Horowitz, B. Z., Prebish, A., Slawson, M., McNair, J., ... Dunn, A. (2018). Notes from the field: Acute poisonings from a synthetic cannabinoid sold as cannabidiol—Utah, 2017–2018. *MMWR. Morbidity and Mortality Weekly Report*, 67(20), 587–588. <https://doi.org/10.15585/mmwr.mm6720a5>
- Huestis, M. A., Gorelick, D. A., Heishman, S. J., Preston, K. L., Nelson, R. A., Moolchan, E. T., & Frank, R. A. (2001). Blockade of effects of smoked marijuana by the CB 1 -selective cannabinoid receptor antagonist SR141716. *Archives of General Psychiatry*, 58(4), 322–328. <https://doi.org/10.1001/archpsyc.58.4.322>
- Huestis, M. A., Solimini, R., Pichini, S., Pacifici, R., Carlier, J., & Busardò, F. P. (2019). Cannabidiol adverse effects and toxicity. *Current Neuropharmacology*, 17(10), 974–989. <https://doi.org/10.2174/1570159X17666190603171901>
- Hurd, Y. L., Spriggs, S., Alishayev, J., Winkel, G., Gurgov, K., Kudrich, C., ... Salsitz, E. (2019). Cannabidiol for the reduction of cue-induced craving and anxiety in drug-abstinent individuals with heroin use disorder: A double-blind randomized placebo-controlled trial. *American Journal of Psychiatry*, 176(11), 911–922. <https://doi.org/10.1176/appi.ajp.2019.18101191>
- Husni, A. S., McCurdy, C. R., Radwan, M. M., Ahmed, S. A., Slade, D., Ross, S. A., ... Cutler, S. J. (2014). Evaluation of phytocannabinoids from high potency Cannabis sativa using in vitro bioassays to determine structure-activity relationships for cannabinoid receptor 1 and cannabinoid receptor 2. *Medicinal Chemistry Research*, 23(9), 4295–4300. <https://doi.org/10.1007/s00044-014-0972-6>
- Ibeas Bih, C., Chen, T., Nunn, A. V., Bazelot, M., Dallas, M., & Whalley, B. J. (2015). Molecular targets of cannabidiol in neurological disorders. *Neurotherapeutics*, 12(4), 699–730. <https://doi.org/10.1007/s13311-015-0377-3>
- Jiang, R., Yamaori, S., Takeda, S., Yamamoto, I., & Watanabe, K. (2011). Identification of cytochrome P450 enzymes responsible for metabolism of cannabidiol by human liver microsomes. *Life Sciences*, 89(5-6), 165–170. <https://doi.org/10.1016/j.lfs.2011.05.018>
- Jikomes, N., & Zoorob, M. (2018). The cannabinoid content of legal cannabis in Washington state varies systematically across testing facilities and popular consumer products. *Scientific Reports*, 8(1), 4519. <https://doi.org/10.1038/s41598-018-22755-2>
- Kaminski, N. E. (1996). Immune regulation by cannabinoid compounds through the inhibition of the cyclic AMP signaling cascade and altered gene expression. *Biochemical Pharmacology*, 52(8), 1133–1140. [https://doi.org/10.1016/0006-2952\(96\)00480-7](https://doi.org/10.1016/0006-2952(96)00480-7)
- Kenne, D. R., Fischbein, R. L., Tan, A. S., & Banks, M. (2017). The use of substances other than nicotine in electronic cigarettes among college students. *Substance Abuse: Research and Treatment*, 11, 1178221817733736. <https://doi.org/10.1177/1178221817733736>

- Kuhathasan, N., Dufort, A., MacKillop, J., Gottschalk, R., Minuzzi, L., & Frey, B. N. (2019). The use of cannabinoids for sleep: A critical review on clinical trials. *Experimental and Clinical Psychopharmacology*, 27(4), 383–401. <https://doi.org/10.1037/pha0000285>
- Lenton, S., Frank, V. A., Barratt, M. J., Potter, G. R., & Decorte, T. (2018). Growing practices and the use of potentially harmful chemical additives among a sample of small-scale cannabis growers in three countries. *Drug and Alcohol Dependence*, 192, 250–256. <https://doi.org/10.1016/j.drugalcdep.2018.07.040>
- Lötsch, J., Weyer-Menkhoff, I., & Tegeder, I. (2018). Current evidence of cannabinoid-based analgesia obtained in preclinical and human experimental settings. *European Journal of Pain (London, England)*, 22(3), 471–484. <https://doi.org/10.1002/ejp.1148>
- Lucas, C. J., Galettis, P., & Schneider, J. (2018). The pharmacokinetics and the pharmacodynamics of cannabinoids. *British Journal of Clinical Pharmacology*, 84(11), 2477–2482. <https://doi.org/10.1111/bcp.13710>
- Mandolini, G. M., Lazzaretti, M., Pigoni, A., Oldani, L., Delvecchio, G., & Brambilla, P. (2018). Pharmacological properties of cannabidiol in the treatment of psychiatric disorders: A critical overview. *Epidemiology and Psychiatric Sciences*, 27(4), 327–335. <https://doi.org/10.1017/S2045796018000239>
- Masataka, N. (2019). Anxiolytic effects of repeated cannabidiol treatment in teenagers with social anxiety disorders. *Frontiers in Psychology*, 10, 2466. <https://doi.org/10.3389/fpsyg.2019.02466>
- McGuire, P., Robson, P., Cubala, W. J., Vasile, D., Morrison, P. D., Barron, R., ... Wright, S. (2018). Cannabidiol (CBD) as an adjunctive therapy in schizophrenia: A multicenter randomized controlled trial. *The American Journal of Psychiatry*, 175(3), 225–231. <https://doi.org/10.1176/appi.ajp.2017.17030325>
- Mead, A. (2019). Legal and regulatory issues governing cannabis and cannabis-derived products in the United States. *Frontiers in Plant Science*, 10, 697. <https://doi.org/10.3389/fpls.2019.00697>
- Mijangos-Moreno, S., Poot-Aké, A., Arankowsky-Sandoval, G., & Murillo-Rodríguez, E. (2014). Intrahypothalamic injection of cannabidiol increases the extracellular levels of adenosine in nucleus accumbens in rats. *Neuroscience Research*, 84, 60–63. <https://doi.org/10.1016/j.neures.2014.04.006>
- Millar, S. A., Stone, N. L., Bellman, Z. D., Yates, A. S., England, T. J., & O'Sullivan, S. E. (2019). A systematic review of cannabidiol dosing in clinical populations. *British Journal of Clinical Pharmacology*, 85(9), 1888–1900. <https://doi.org/10.1111/bcp.14038>
- Morrison, G., Crockett, J., Blakey, G., & Sommerville, K. (2019). A phase 1, open-label, pharmacokinetic trial to investigate possible drug-drug interactions between clobazam, stiripentol, or valproate and cannabidiol in healthy subjects. *Clinical Pharmacology in Drug Development*, 8(8), 1009–1031. <https://doi.org/10.1002/cpdd.665>
- Munnangi, S., & Angus, L. D. (2019). *Placebo effect*. StatPearls. Treasure Island, FL: StatPearls Publishing.
- O'Connell, B. K., Gloss, D., & Devinsky, O. (2017). Cannabinoids in treatment-resistant epilepsy: A review. *Epilepsy & Behavior*, 70(Pt B), 341–348. <https://doi.org/10.1016/j.yebeh.2016.11.012>
- O'Connor, S. M., & Lietzan, E. (2019). The surprising reach of FDA regulation of cannabis, even after descheduling. *The American University Law Review*, 68(3), 823–925.
- Ohlsson, A., Lindgren, J. E., Andersson, S., Agurell, S., Gillespie, H., & Hollister, L. E. (1986). Single-dose kinetics of deuterium-labelled cannabidiol in man after smoking and intravenous administration. *Biomedical & Environmental Mass Spectrometry*, 13(2), 77–83. <https://doi.org/10.1002/bms.1200130206>
- Pate, D. W. (1994). Chemical ecology of cannabis. *Journal of the International Hemp Association*, 29, 32–37.
- Pavlovic, R., Nenna, G., Calvi, L., Panseri, S., Borgonovo, G., Giupponi, L., ... Giorgi, A. (2018). Quality traits of “cannabidiol oils”: Cannabinoids content, terpene fingerprint and oxidation stability of European commercially available preparations. *Molecules (Basel, Switzerland)*, 23(5), 1230. <https://doi.org/10.3390/molecules23051230>
- Pellati, F., Borgonetti, V., Brighenti, V., Biagi, M., Benvenuti, S., & Corsi, L. (2018). *Cannabis sativa* L. and nonpsychoactive cannabinoids: Their chemistry and role against oxidative stress, inflammation, and cancer. *BioMed Research International*, 2018, 1691428. <https://doi.org/10.1155/2018/1691428>
- Pollan, M. (2001). *The botany of desire: A plant's-eye view of the world*. New York: Random House.

- Pound, P., & Ritskes-Hoitinga, M. (2018). Is it possible to overcome issues of external validity in preclinical animal research? Why most animal models are bound to fail. *Journal of Translational Medicine*, 16(1), 304. <https://doi.org/10.1186/s12967-018-1678-1>
- Premoli, M., Aria, F., Bonini, S. A., Maccarinelli, G., Gianoncelli, A., Pina, S. D., ... Mastinu, A. (2019). Cannabidiol: Recent advances and new insights for neuropsychiatric disorders treatment. *Life Sciences*, 224, 120–127. <https://doi.org/10.1016/j.lfs.2019.03.053>
- Raber, J. C., Elzinga, S., & Kaplan, C. (2015). Understanding dabs: Contamination concerns of cannabis concentrates and cannabinoid transfer during the act of dabbing. *The Journal of Toxicological Sciences*, 40(6), 797–803. <https://doi.org/10.2131/jts.40.797>
- Rahn, B. (2018). Which cannabis strains are highest in CBD, according to lab data? <https://www.leafly.com/news/cbd/high-cbd-marijuana-strains-according-to-lab-data>
- Riebe, C. J., & Wotjak, C. T. (2011). Endocannabinoids and stress. *Stress (Amsterdam, Netherlands)*, 14(4), 384–397. <https://doi.org/10.3109/10253890.2011.586753>
- Roberts, B. A. (2019). Legalized cannabis in Colorado emergency departments: A cautionary review of negative health and safety effects. *The Western Journal of Emergency Medicine*, 20(4), 557–572. <https://doi.org/10.5811/westjem.2019.4.39935>
- Rock, E. M., Bolognini, D., Limebeer, C. L., Cascio, M. G., Anavi-Goffer, S., Fletcher, P. J., ... Parker, L. A. (2012). Cannabidiol, a non-psychotropic component of cannabis, attenuates vomiting and nausea-like behaviour via indirect agonism of 5-HT_{1A} somatodendritic autoreceptors in the dorsal raphe nucleus. *British Journal of Pharmacology*, 165(8), 2620–2634. <https://doi.org/10.1111/j.1476-5381.2011.01621.x>
- Rodríguez de Fonseca, F., Del Arco, I., Bermudez-Silva, F. J., Bilbao, A., Cippitelli, A., & Navarro, M. (2005). The endocannabinoid system: Physiology and pharmacology. *Alcohol and Alcoholism (Oxford, Oxfordshire)*, 40(1), 2–14. <https://doi.org/10.1093/alcalc/agh110>
- Romano, L. L., & Hazekamp, A. (2013). Cannabis oil: Chemical evaluation of an upcoming cannabis-based medicine. *Cannabinoids*, 1(1), 1–11.
- Rubin, R. (2018). The path to the first FDA-approved cannabis-derived treatment and what comes next. *Journal of the American Medical Association*, 320(12), 1227–1229. <https://doi.org/10.1001/jama.2018.11914>
- Ruiz de Azua, I., & Lutz, B. (2019). Multiple endocannabinoid-mediated mechanisms in the regulation of energy homeostasis in brain and peripheral tissues. *Cellular and Molecular Life Sciences*, 76(7), 1341–1363. <https://doi.org/10.1007/s00018-018-2994-6>
- Russo, E. B. (2011). Taming THC: Potential cannabis synergy and phytocannabinoid-terpenoid entourage effects. *British Journal of Pharmacology*, 163(7), 1344–1364. <https://doi.org/10.1111/j.1476-5381.2011.01238.x>
- Sagredo, O., Ramos, J. A., Decio, A., Mechoulam, R., & Fernández-Ruiz, J. (2007). Cannabidiol reduced the striatal atrophy caused 3-nitropropionic acid in vivo by mechanisms independent of the activation of cannabinoid, vanilloid TRPV1 and adenosine A_{2A} receptors. *The European Journal of Neuroscience*, 26(4), 843–851. <https://doi.org/10.1111/j.1460-9568.2007.05717.x>
- Schoedel, K. A., Szeto, I., Setnik, B., Sellers, E. M., Levy-Cooperman, N., Mills, C., ... Sommerville, K. (2018). Abuse potential assessment of cannabidiol (CBD) in recreational polydrug users: A randomized, double-blind, controlled trial. *Epilepsy & Behavior*, 88, 162–171. <https://doi.org/10.1016/j.yebeh.2018.07.027>
- Seltenrich, N. (2019). Into the weeds: Regulating pesticides in cannabis. *Environmental Health Perspectives*, 127(4), 042001. <https://doi.org/10.1289/EHP5265>
- Shannon, S., Lewis, N., Lee, H., & Hughes, S. (2019). Cannabidiol in anxiety and sleep: A large case series. *The Permanente Journal*, 23, 18–041. <https://doi.org/10.7812/TPP/18-041>
- Siano, F., Moccia, S., Picariello, G., Russo, G. L., Sorrentino, G., Di Stasio, M., ... Volpe, M. G. (2018). Comparative study of chemical, biochemical characteristic and ATR-FTIR analysis of seeds, oil and flour of the edible Fedora cultivar hemp (*Cannabis sativa* L.). *Molecules (Basel, Switzerland)*, 24(1), 83. <https://doi.org/10.3390/molecules24010083>
- Smart, R., Caulkins, J. P., Kilmer, B., Davenport, S., & Midgette, G. (2017). Variation in cannabis potency and prices in a newly legal market: Evidence from 30 million cannabis sales in Washington state. *Addiction (Abingdon, England)*, 112(12), 2167–2177. <https://doi.org/10.1111/add.13886>

- Sonego, A. B., Gomes, F. V., Del Bel, E. A., & Guimaraes, F. S. (2016). Cannabidiol attenuates haloperidol-induced catalepsy and c-Fos protein expression in the dorsolateral striatum via 5-HT1A receptors in mice. *Behavioural Brain Research*, *309*, 22–28. <https://doi.org/10.1016/j.bbr.2016.04.042>
- Stogner, J. M., & Miller, B. L. (2015). Assessing the dangers of “dabbing”: Mere marijuana or harmful new trend? *Pediatrics*, *136*(1), 1–3. <https://doi.org/10.1542/peds.2015-0454>
- Szaflarski, J. P., Bebin, E. M., Cutter, G., DeWolfe, J., Dure, L. S., Gaston, T. E., ... Ver Hoef, L. W., & the UAB CBD Program. (2018). Cannabidiol improves frequency and severity of seizures and reduces adverse events in an open-label add-on prospective study. *Epilepsy & Behavior*, *87*, 131–136. <https://doi.org/10.1016/j.yebeh.2018.07.020>
- Taylor, L., Gidal, B., Blakey, G., Tayo, B., & Morrison, G. (2018). A phase I, randomized, double-blind, placebo-controlled, single ascending dose, multiple dose, and food effect trial of the safety, tolerability and pharmacokinetics of highly purified cannabidiol in healthy subjects. *CNS Drugs*, *32*(11), 1053–1067. <https://doi.org/10.1007/s40263-018-0578-5>
- Tham, M., Yilmaz, O., Alaverdashvili, M., Kelly, M. E. M., Denovan-Wright, E. M., & Laprairie, R. B. (2019). Allosteric and orthosteric pharmacology of cannabidiol and cannabidiol-dimethylheptyl at the type 1 and type 2 cannabinoid receptors. *British Journal of Pharmacology*, *176*(10), 1455–1469. <https://doi.org/10.1111/bph.14440>
- Thomas, B. F., & ElSohly, M. A. (2015). *The analytical chemistry of cannabis: Quality assessment, assurance, and regulation of medicinal marijuana and cannabinoid preparations (Vol. 1)*. Cambridge, MA: Elsevier in cooperation with RTI Press.
- US Food and Drug Administration. (2018). *FDA approves first drug comprised of an active ingredient derived from marijuana to treat rare, severe forms of epilepsy* [Press release]. Retrieved from <https://www.fda.gov/news-events/press-announcements/fda-approves-first-drug-comprised-active-ingredient-derived-marijuana-treat-rare-severe-forms>
- US Department of Agriculture. (2019, June 26). *USDA Farm Bill implementation progress update*. Retrieved from <https://www.usda.gov/media/press-releases/2019/06/26/usda-farm-bill-implementation-progress-update>
- US Drug Enforcement Administration. (n.d.). *Drug scheduling*. Retrieved from <https://www.dea.gov/drug-scheduling>
- VanDolah, H. J., Bauer, B. A., & Mauck, K. F. (2019). Clinicians’ guide to cannabidiol and hemp oils. *Mayo Clinic Proceedings*, *94*(9), 1840–1851. <https://doi.org/10.1016/j.mayocp.2019.01.003>
- Vandrey, R., Raber, J. C., Raber, M. E., Douglass, B., Miller, C., & Bonn-Miller, M. O. (2015). Cannabinoid dose and label accuracy in edible medical cannabis products. *Journal of the American Medical Association*, *313*(24), 2491–2493. <https://doi.org/10.1001/jama.2015.6613>
- Ward, S. J., Ramirez, M. D., Neelakantan, H., & Walker, E. A. (2011). Cannabidiol prevents the development of cold and mechanical allodynia in paclitaxel-treated female C57Bl6 mice. *Anesthesia and Analgesia*, *113*(4), 947–950. <https://doi.org/10.1213/ANE.0b013e3182283486>
- White, C. M. (2019). A review of human studies assessing cannabidiol’s (CBD) therapeutic actions and potential. *Journal of Clinical Pharmacology*, *59*(7), 923–934. <https://doi.org/10.1002/jcph.1387>
- Wiley, J. L., Lowe, J. A., Balster, R. L., & Martin, B. R. (1995). Antagonism of the discriminative stimulus effects of delta 9-tetrahydrocannabinol in rats and rhesus monkeys. *Journal of Pharmacology and Experimental Therapeutics*, *275*(1), 1–6.
- Wilkinson, J. D., Whalley, B. J., Baker, D., Pryce, G., Constanti, A., Gibbons, S., & Williamson, E. M. (2003). Medicinal cannabis: Is delta9-tetrahydrocannabinol necessary for all its effects? *Journal of Pharmacy and Pharmacology*, *55*(12), 1687–1694. <https://doi.org/10.1211/0022357022304>
- Zuardi, A. W. (2006). History of cannabis as a medicine: A review. *Brazilian Journal of Psychiatry*, *28*(2), 153–157. <https://doi.org/10.1590/S1516-44462006000200015>

RTI International is an independent, nonprofit research institute dedicated to improving the human condition. We combine scientific rigor and technical expertise in social and laboratory sciences, engineering, and international development to deliver solutions to the critical needs of clients worldwide.

www.rti.org/rtipress

RTI Press publication OP-0065-2004